

APPLICATION UNDER UNITED STATES PATENT LAWS

Atty. Dkt. No. PW 278416
(M#)

Invention: L-LYSINE-PRODUCING CORYNEBACTERIA AND PROCESS FOR THE PREPARATION OF LYSINE

Inventor (s): Caroline KREUTZER
Bettina MÖCKEL
Walter PFEFFERLE
Lothar EGDELING
Hermann SAHM
Miroslav PATEK

Pillsbury Winthrop LLP
Intellectual Property Group
1100 New York Avenue, NW
Ninth Floor
Washington, DC 20005-3918
Attorneys
Telephone: (202) 861-3000

This is a:

- ☐ Provisional Application
- ☐ Regular Utility Application
- ☒ Continuation-In-Part Application
 - ☒ The contents of the parent are incorporated by reference
- ☐ PCT National Phase Application
- ☐ Design Application
- ☐ Reissue Application
- ☐ Plant Application
- ☐ Substitute Specification
 - Sub. Spec Filed _____
 - in App. No. _____ / _____
- ☐ Marked up Specification re
 - Sub. Spec. filed _____
 - In App. No _____ / _____

SPECIFICATION

L-Lysine-Producing Corynebacteria and Process for the Preparation of Lysine

This application is a continuation-in-part of U.S.

- 5 application no. 09/353,608, filed July 14, 1999, which is incorporated herein by reference in its entirety.

BACKGROUND OF THE INVENTION

Field of the Invention

- 10 The invention relates to L-lysine-producing strains of corynebacteria with enhanced pyc gene (pyruvate carboxylase gene), in which strains additional genes, chosen from the group consisting of the dapA gene (dihydrodipicolinate synthase gene), the lysC gene
- 15 (aspartate kinase gene), the lysE gene (lysine export carrier gene) and the dapB gene (dihydrodipicolinate reductase gene), but especially the dapA gene, are enhanced and, in particular, over-expressed, and to a process for the preparation of L-lysine.

20 Background Information

L-Lysine is a commercially important L-amino acid which is used especially as a feed additive in animal nutrition. The demand of such feed additives has been steadily increasing in recent years.

- 25 L-Lysine is prepared by a fermentation process with L-lysine-producing strains of corynebacteria, especially *Corynebacterium glutamicum*. Because of the great importance of this product, attempts are constantly being made to improve the preparative process. Improvements to
- 30 the process may relate to measures involving the fermentation technology, e.g. stirring and oxygen supply, or the composition of the nutrient media, e.g. the sugar concentration during fermentation, or the work-up to the product form, e.g. by ion exchange chromatography, or the

intrinsic productivity characteristics of the microorganism itself.

The productivity characteristics of these microorganisms are improved by using methods of mutagenesis, selection
5 and mutant choice to give strains which are resistant to antimetabolites, e.g. S-(2-aminoethyl)cysteine, or auxotrophic for amino acids, e.g. L-leucine, and produce L-lysine.

Methods of recombinant DNA technology have also been used
10 for some years in order to improve L-lysine-producing strains of *Corynebacterium glutamicum* by amplifying individual biosynthesis genes and studying the effect on the L-lysine production.

Thus, EP-A-0 088 166 reports the increase in productivity,
15 after amplification, of a DNA fragment conferring resistance to aminoethylcysteine. EP-B-0 387 527 reports the increase in productivity, after amplification, of an lysC allele coding for a feedback-resistant aspartate kinase. EP-B-0 197 335 reports the increase in
20 productivity, after amplification, of the dapA gene coding for dihydrodipicolinate synthase. EP-A-0 219 027 reports the increase in productivity, after amplification, of the asd gene coding for aspartate semialdehyde dehydrogenase. Pisabarro et al. (Journal of Bacteriology 175(9), 2743-
25 2749 (1993)) describe the dapB gene coding for dihydrodipicolinate reductase.

The effect of the amplification of primary metabolism genes on the L-lysine production has also been studied. Thus EP-A-0 219 027 reports the increase in productivity,
30 after amplification, of the aspC gene coding for aspartate aminotransferase. EP-B-0 143 195 and EP-B-0 358 940 report the increase in productivity, after amplification, of the ppc gene coding for phosphoenolpyruvate carboxylase. DE-A-198 31 609 reports the increase in

productivity, after amplification, of the *pyc* gene coding for pyruvate carboxylase. The anaplerotic reaction catalyzed by pyruvate carboxylase is of particular importance compared with the reaction catalyzed by phosphoenolpyruvate carboxylase. Thus, Wendisch et al. (FEMS Microbiology Letters 112, 269-274 (1993)) showed that the lysine production of the strain MH20-22B was not impaired by turning off the *ppc* gene.

Finally, DE-A-195 48 222 describes that an increased activity of the L-lysine export carrier coded for by the *lysE* gene promotes lysine production.

In addition to these attempts to amplify an individual gene, attempts have also been made to amplify two or more genes simultaneously, thereby improving the L-lysine production in corynebacteria. Thus, DE-A-38 23 451 reports the increase in productivity, after simultaneous amplification, of the *asd* gene and the *dapA* gene from *Escherichia coli*. DE-A-39 43 117 discloses the increase in productivity, after simultaneous amplification, of an *lysC* allele coding for a feedback-resistant aspartate kinase and of the *dapA* gene. EP-A-0 841 395 particularly reports the increase in productivity, after simultaneous amplification, of an *lysC* allele coding for a feedback-resistant aspartate kinase and of the *dapB* gene; further improvements could be achieved by additional amplification of the *dapB*, *lysA* and *ddh* genes. EP-A-0 854 189 describes the increase in productivity, after simultaneous amplification, of an *lysC* allele coding for a feedback-resistant aspartate kinase and of the *dapA*, *dapB*, *lysA* and *aspC* genes. EP-A-0 857 784 particularly reports the increase in productivity, after simultaneous amplification, of an *lysC* allele coding for a feedback-resistant aspartate kinase enzyme and the *lysA* gene; a further improvement could be achieved by additional amplification of the *ppc* gene.

It is clear from the many processes described in the state of the art, that there is a need for the development of novel approaches and for the improvement of existing processes for lysine production with corynebacteria.

SUMMARY OF THE INVENTION

Object of the Invention

It is an object of the invention to provide novel L-lysine-producing strains of corynebacteria and processes
5 for the preparation of L-lysine.

Description of the Invention

When L-lysine or lysine is mentioned in the following text, it should be understood as meaning not only the base but also the appropriate salts, e.g. lysine hydrochloride
10 or lysine sulfate.

The invention provides L-lysine-producing strains of corynebacteria with enhanced pyc gene (pyruvate carboxylase gene), wherein additional genes, chosen from the group consisting of the dapA gene (dihydrodipicolinate synthase gene), the lysC gene (aspartate kinase gene), the
15 lysE gene (lysine export carrier gene) and the dapB gene (dihydrodipicolinate reductase gene), but especially the dapA gene, are enhanced and, in particular, over-expressed.

20 A novel DNA sequence located upstream (5' end) from the dapB gene has also been found, which carries the -35 region of the dapB promoter and is advantageous for the expression of the dapB gene. It is shown as SEQ ID No. 1.

A corresponding DNA capable of replication, with the
25 nucleotide sequence shown in SEQ ID No. 1, is also included in the invention.

The invention also provides the MC20 and MA16 mutations of the dapA promoter shown in SEQ ID No. 5 and SEQ ID No. 6, deposited in the strains DSM12868 and DSM12867.

30 The invention furthermore provides L-lysine-producing strains of corynebacteria with amplified pyc gene, wherein

additionally the dapA and dapB genes are simultaneously enhanced and, in particular, over-expressed.

The invention also provides L-lysine-producing strains of corynebacteria with enhanced pyc gene, wherein
5 additionally the dapA, dapB and lysE genes are simultaneously enhanced and, in particular, over-expressed.

In this context the term "enhancement" describes the increase in the intracellular activity, in a
10 microorganism, of one or more enzymes which are coded for by the appropriate DNA, by increasing the copy number of the gene(s), using a strong promoter or using a gene coding for an appropriate enzyme with a high activity, and optionally combining these measures.

15 In this context, "amplification" describes a specific procedure for achieving an enhancement whereby the number of DNA molecules carrying a gene or genes, an allele or alleles, a regulatory signal or signals or any other genetic feature(s) is increased.

20 The invention also provides a process for the preparation of L-lysine using the bacteria described above.

The microorganisms which the present invention provides can prepare L-lysine from glucose, sucrose, lactose, fructose, maltose, molasses, starch or cellulose or from
25 glycerol and ethanol, especially from glucose or sucrose. Said microorganisms are corynebacteria, especially of the genus Corynebacterium. The species Corynebacterium glutamicum may be mentioned in particular in the genus Corynebacterium, being known to those skilled in the art
30 for its ability to produce amino acids. This species includes wild-type strains such as Corynebacterium glutamicum ATCC13032, Brevibacterium flavum ATCC14067, Corynebacterium melassecola ATCC17965 and strains or

mutants derived therefrom. Examples of L-lysine-producing mutants of corynebacteria are:

Corynebacterium glutamicum FERM-P 1709
Brevibacterium flavum FERM-P 1708
5 Brevibacterium lactofermentum FERM-P 1712
Brevibacterium flavum FERM-P 6463
Brevibacterium flavum FERM-P 6464
Corynebacterium glutamicum DSM5714
Corynebacterium glutamicum DSM12866

10 It has now been found that an enhanced expression of the
lysE gene in addition to the pyc gene, or an additionally
enhanced expression of an lysC allele coding for a
feedback-resistant aspartate kinase, or an additionally
enhanced expression of the dapB gene and, in particular,
15 an additionally enhanced expression of the dapA gene,
individually or together, further improve L-lysine
production.

It has also been found that, for a given over-expression
of the pyc gene, the simultaneous, additionally enhanced
20 expression of the dapA and dapB genes brings further
advantages for L-lysine production.

Finally, it has been found that, for a given over-
expression of the pyc gene, the simultaneous, additionally
enhanced expression of the dapA, dapB and lysE genes is
25 extremely advantageous for L-lysine production.

An enhancement (over-expression) is achieved, e.g., by
increasing the copy number of the appropriate genes or
mutating the promoter and regulatory region or the
ribosome binding site located upstream from the structural
30 gene. Expression cassettes incorporated upstream from the
structural gene function in the same way. Inducible
promoters additionally make it possible to increase the
expression in the course of the formation of L-lysine by
fermentation. Measures for prolonging the life of the

mRNA also improve the expression. Furthermore, the enzyme activity is also enhanced by preventing the degradation of the enzyme protein, the genes or gene constructs either being located in plasmids (shuttle vectors) of variable
5 copy number or being integrated and amplified in the chromosome. Alternatively, it is also possible to achieve an over-expression of the genes in question by changing the composition of the media and the culture technique.

Those skilled in the art will find relevant instructions
10 inter alia in Martin et al. (Bio/Technology 5, 137-146 (1987)), Guerrero et al. (Gene 138, 35-41 (1994)), Tsuchiya and Morinaga (Bio/Technology 6, 428-430 (1988)), Eikmanns et al. (Gene 102, 93-98 (1991)), EP-0 472 869, US 4,601,893, Schwarzer and Pühler (Bio/Technology 9, 84-87
15 (1991)), Reinscheid et al. (Applied and Environmental Microbiology 60, 126-132 (1994)), LaBarre et al. (Journal of Bacteriology 175, 1001-1007 (1993)), WO 96/15246, Malumbres et al. (Gene 134, 15-24 (1993)), JP-A-10-229891, Jensen and Hammer (Biotechnology and Bioengineering 58,
20 191-195 (1998)) or the handbook "Manual of Methods for General Bacteriology" of the American Society for Bacteriology (Washington DC, USA, 1981) and well-known textbooks on genetics and molecular biology.

The genes from *Corynebacterium glutamicum* used according
25 to the invention are described and can be isolated, prepared or synthesized by known methods.

Methods of localized mutagenesis are described, *inter alia*, by Higuchi et al. (Nucleic Acids Research 16, 7351-7367 (1988)) or by Silver et al. in the handbook by Innis,
30 Glefand and Sninsky (eds.) entitled "PCR Strategies" (Academic Press, London, UK, 1995).

The first step in isolating a gene of interest from *C. glutamicum* is to construct a gene library of this microorganism in, e.g., *E. coli* or optionally also in *C.*

glutamicum. The construction of gene libraries is documented in generally well-known textbooks and handbooks. Examples which may be mentioned are the textbook by Winnacker entitled "From Genes to Clones, Introduction to Gene Technology" (Verlag Chemie, Weinheim, Germany, 1990) or the handbook by Sambrook et al. entitled "Molecular Cloning, A Laboratory Manual" (Cold Spring Harbor Laboratory Press, 1989). Bathe et al. (Molecular and General Genetics, 252: 255-265 (1996)) describe a gene library of *C. glutamicum* ATCC13032 which was constructed using cosmid vector SuperCos I (Wahl et al., Proceedings of the National Academy of Sciences, USA 84, 2160-2164 (1987)) in *E. coli* K-12 NM554 (Raleigh et al., Nucleic Acids Research 16: 1563-1575 (1988)). Börmann et al. (Molecular Microbiology 6(3), 317-326) in turn describe a gene library of *C. glutamicum* ATCC13032 using cosmid pHC79 (Hohn and Collins, Gene 11, 291-298 (1980)). A gene library of *C. glutamicum* in *E. coli* can also be constructed using plasmids like pBR322 (Bolivar, Life Sciences 25, 807-818 (1979)) or pUC19 (Norrandar et al., Gene 26: 101-106 (1983)). In the same way, it is also possible to use shuttle vectors such as pJC1 (Cremer et al., Molecular and General Genetics 220, 478-480 (1990)) or pEC5 (Eikmanns et al., Gene 102, 93-98 (1991)), which replicate in *E. coli* and *C. glutamicum*. Restriction- and/or recombination-defective strains are particularly suitable hosts, an example being the *E. coli* strain DH5 α mcr, which has been described by Grant et al. (Proceedings of the National Academy of Sciences, USA 87, 4645-4649 (1990)). Other examples are the restriction-defective *C. glutamicum* strains RM3 and RM4, which are described by Schäfer et al. (Applied and Environmental Microbiology 60(2), 756-759 (1994)).

The gene library is then transferred to an indicator strain by transformation (Hanahan, Journal of Molecular Biology 166, 557-580 (1983)) or electroporation (Tauch et

al., FEMS Microbiological Letters, 123: 343-347 (1994)).

The characteristic feature of the indicator strain is that it possesses a mutation in the gene of interest which causes a detectable phenotype, e.g. an auxotrophy. The indicator strains or mutants are obtainable from publicized sources or strain collections, e.g. the Genetic Stock Center of Yale University (New Haven, Connecticut, USA), or if necessary are specially prepared. An example of such an indicator strain which may be mentioned is the E. coli strain RDA8 requiring mesodiaminopimelic acid (Richaud et al., C.R. Acad. Sci. Paris Ser. III 293: 507-512 (1981)), which carries a mutation (dapA::Mu) in the dapA gene.

After transformation of the indicator strain with a recombinant plasmid carrying the gene of interest, and expression of the gene in question, the indicator strain becomes prototrophic in respect of the appropriate characteristic. If the cloned DNA fragment confers resistance, e.g. to an antimetabolite such as S-(2-aminoethyl)cysteine, the indicator strain carrying the recombinant plasmid can be identified by selection on appropriately supplemented nutrient media.

If the nucleotide sequence of the gene region of interest is known or obtainable from a data bank, the chromosomal DNA can be isolated by known methods, e.g. as described by Eikmanns et al. (Microbiology 140, 1817-1828 (1994)), and the gene in question can be synthesized by the polymerase chain reaction (PCR) using suitable primers and cloned into a suitable plasmid vector, e.g. pCRIITOP0 from Invitrogen (Groningen, The Netherlands). A summary of PCR methodology can be found in the book by Newton and Graham entitled "PCR" (Spektrum Akademischer Verlag, Heidelberg, Germany, 1994).

Examples of publicly accessible data banks for nucleotide sequences are that of the European Molecular Biologies

Laboratories (EMBL, Heidelberg, Germany) or that of the National Center for Biotechnology Information (NCBI, Bethesda, MD, USA).

The isolation and cloning of the *pyc* gene from *C.*

- 5 *glutamicum* ATCC13032 are described in DE-A-198 31 609 and by Koffas et al. (Applied Microbiology and Biotechnology 50, 346-352 (1988)). The nucleotide sequence of the *pyc* gene is obtainable under accession number AF038548 or Y09548.

- 10 The isolation and cloning of the *lysE* gene from *C. glutamicum* ATCC13032 are described in DE-A-195 48 222. The nucleotide sequence of the *lysE* gene is obtainable under accession number X96471.

- The isolation, cloning and sequencing of the *dapA* gene from various strains of *C. glutamicum* are described by
15 Cremer et al. (Molecular and General Genetics 220: 478-480 (1990)), by Pisabarro et al. (Journal of Bacteriology 175: 2743-2749 (1993)) and by Bonnassie et al. (Nucleic Acids Research 18: 6421 (1990)). DE-A-39 43 117 reports the
20 amplification of the *dapA* gene by means of plasmid pJC23. The nucleotide sequence of the *dapA* gene is obtainable under accession number X53993.

- The isolation, cloning and sequencing of the *dapB* gene from *Brevibacterium lactofermentum* are described by
25 Pisabarro et al. (Journal of Bacteriology 175: 2743-2749 (1993)). The nucleotide sequence of the *dapB* gene is obtainable under accession number X67737.

- The isolation, cloning and sequencing of the *lysC* gene and of *lysC* alleles coding for a feedback-resistant aspartate
30 kinase are reported by several authors. Thus, Kalinowski et al. (Molecular and General Genetics 224: 317-324 (1990)) report the *lysC* allele from the *C. glutamicum* strain DM58-1. DE-A-39 43 117 reports the cloning of the *lysC* allele from the *C. glutamicum* strain MH20. Follettie

et al. (Journal of Bacteriology 175: 4096-4103 (1993)) report the lysC allele from the C. flavum strain N13, which is called "ask" in said publication. The nucleotide sequences of the lysC gene and of various lysC alleles are obtainable inter alia under accession numbers X57226 and E06826.

The genes obtained in this way can then be incorporated inter alia into plasmid vectors, e.g. pJC1 (Cremer et al., Molecular and General Genetics 220, 478-480 (1990)) or pEC5 (Eikmanns et al., Gene, 102, 93-98 (1991)), individually or in suitable combinations, transferred to desired strains of corynebacteria, e.g. the strain MH20-22B (Schrumpf et al., Applied Microbiology and Biotechnology 37: 566-571 (1992)), by transformation, e.g. as in Thierbach et al. (Applied Microbiology and Biotechnology 29, 356-362 (1988)), or by electroporation, e.g. as in Dunican and Shivnan (Bio/Technology 7, 1067-1070 (1989)), and expressed. The strain to be chosen can equally well be transformed with two plasmid vectors, each containing the gene or genes in question, thereby achieving the advantageous, simultaneously enhanced expression of two or more genes in addition to the known enhancement of the pyc gene.

Examples of such strains are:

- the strain MH20-22B/pJC23/pEC7pyc, in which the pyc and dapA genes are simultaneously enhanced, or
- the strain MH20-22B/pJC33/pEC7pyc, in which the pyc- and the lysC(FBR) allele are simultaneously enhanced and, in particular, over-expressed, or
- the strain MH20-22B/pJC23/pEC7dapBpyc, in which the pyc, dapA and dapB genes are simultaneously enhanced and, in particular, over-expressed, or

- the strain MH20-22B/pJC23/pEC7lysEdapBpyc, in which the pyc, dapA, dapB and lysE genes are simultaneously enhanced and, in particular, over-expressed.

The microorganisms prepared according to the invention can
5 be cultivated for L-lysine production continuously or
discontinuously by the batch process, the fed batch
process or the repeated fed batch process. A summary of
known cultivation methods is provided in the textbook by
Chmiel (Bioprozesstechnik 1. Einführung in die
10 Bioverfahrenstechnik (Bioprocess Technology 1.
Introduction to Bioengineering) (Gustav Fischer Verlag,
Stuttgart, 1991)) or in the textbook by Storhas
(Bioreaktoren und periphere Einrichtungen (Bioreactors and
Peripheral Equipment) (Vieweg Verlag, Brunswick/Wiesbaden,
15 1994)).

The culture medium to be used must appropriately meet the
demands of the particular microorganisms. Descriptions of
culture media for various microorganisms can be found in
the handbook "Manual of Methods for General Bacteriology"
20 of the American Society for Bacteriology (Washington D.C.,
USA, 1981). Carbon sources which can be used are sugars
and carbohydrates, e.g. glucose, sucrose, lactose,
fructose, maltose, molasses, starch and cellulose, oils
and fats, e.g. soya oil, sunflower oil, groundnut oil and
25 coconut fat, fatty acids, e.g. palmitic acid, stearic acid
and linoleic acid, alcohols, e.g. glycerol and ethanol,
and organic acids, e.g. acetic acid. These substances can
be used individually or as a mixture. Nitrogen sources
which can be used are organic nitrogen-containing
30 compounds such as peptones, yeast extract, meat extract,
malt extract, corn steep liquor, soybean flour and urea,
or inorganic compounds such as ammonium sulfate, ammonium
chloride, ammonium phosphate, ammonium carbonate and
ammonium nitrate. The nitrogen sources can be used
35 individually or as a mixture. Phosphorus sources which
can be used are potassium dihydrogenphosphate or

dipotassium hydrogenphosphate or the corresponding sodium salts. The culture medium must also contain metal salts, e.g. magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth-promoting substances such as amino acids and vitamins can be used in addition to the substances mentioned above. Said feed materials can be added to the culture all at once or fed in appropriately during cultivation.

The pH of the culture is controlled by the appropriate use of basic compounds such as sodium hydroxide, potassium hydroxide or ammonia, or acid compounds such as phosphoric acid or sulfuric acid. Foaming can be controlled using antifoams such as fatty acid polyglycol esters. The stability of plasmids can be maintained by adding suitable selectively acting substances, e.g. antibiotics, to the medium. Aerobic conditions are maintained by introducing oxygen or oxygen-containing gaseous mixtures, e.g. air, into the culture. The temperature of the culture is normally 20°C to 45°C and preferably 25°C to 40°C. The culture is continued until L-lysine formation has reached a maximum. This objective is normally achieved within 10 hours to 160 hours.

The concentration of L-lysine formed can be determined with the aid of amino acid analyzers by means of ion exchange chromatography and postcolumn reaction with ninhydrin detection, as described by Spackmann et al. (Analytical Chemistry 30, 1190 (1958)).

The following microorganisms have been deposited at the Deutsche Sammlung für Mikroorganismen und Zellkulturen (German Collection of Microorganisms and Cell Cultures (DSMZ), Brunswick, Germany) under the terms of the Budapest Treaty:

- *Escherichia coli* K-12 strain DH5 α /pEC7pyc as DSM12870

- Escherichia coli K-12 strain DH5 α /pEC7dapBpyc as DSM12873
- Escherichia coli K-12 strain DH5 α /pEC7lysEdapBpyc as DSM12874
- 5 • Corynebacterium glutamicum strain DSM5715/pJC23 as DSM12869
- Corynebacterium glutamicum strain DSM5715aecD::dapA(MA16) as DSM12867
- 10 • Corynebacterium glutamicum strain DSM5715aecD::dapA(MC20) as DSM12868
- Corynebacterium glutamicum strain DM678 as DSM12866
- Escherichia coli K-12 strain DH5 α /pEC7lysEpyc as DSM 12872
- Escherichia coli K-12 strain DH5 α /pEC7dapBlysE as DSM 12875
- 15 • Escherichia coli K-12 strain DH5 α /pEC7lysE as DSM 12871

DETAILED DESCRIPTION OF THE INVENTION

The present invention is illustrated in greater detail
20 below with the aid of Examples.

Example 1

Preparation of the DNA coding for the lyse gene

Chromosomal DNA was isolated from the strain ATCC13032 by
the conventional methods (Eikmanns et al., Microbiology
25 140: 1817-1828 (1994)). The polymerase chain reaction
(PCR) was used to amplify a DNA fragment carrying the lyse
gene. The following primer oligonucleotides were chosen
for the PCR on the basis of the lyse gene sequence known

for *C. glutamicum* (Vrljic et al., Molecular Microbiology 22(5), 815-826 (1996)) (accession number X96471):

LysBam1:

5' CTC GAG AGC (GGA TCC) GCG CTG ACT CAC C 3'

5 LysBam2:

5' GGA GAG TAC GGC (GGA TCC) ACC GTG ACC 3'

The primers shown were synthesized by MWG Biotech (Ebersberg, Germany) and the PCR was carried out by the standard PCR method of Innis et al. (PCR Protocols. A Guide to Methods and Applications, 1990, Academic Press). The primers make it possible to amplify an approx. 1.1 kb DNA fragment carrying the *lyse* gene. The primers also contain the sequence for the cleavage site of the restriction endonuclease BamHI, which is indicated by brackets in the nucleotide sequence shown above.

The amplified DNA fragment of approx. 1.1 kb, carrying the *lyse* gene, was identified by means of electrophoresis in 0.8% agarose gel, isolated from the gel and purified with the QIAquick Gel Extraction Kit (cat. no. 28704) from Qiagen (Hilden, Germany).

The fragment was then ligated by means of T4 DNA ligase from Boehringer Mannheim (Mannheim, Germany) to vector pUC18 (Norrandar et al., Gene (26) 101-106 (1983)). This was done by fully cleaving vector pUC18 with the restriction endonuclease SmaI and treating it with alkaline phosphatase (Boehringer Mannheim, Mannheim, Germany). The ligation mixture was transformed to the *E. coli* strain DH5 α (Hanahan, in: DNA Cloning. A Practical Approach, Vol. I, IRL-Press, Oxford, Washington DC, USA). Plasmid-carrying cells were selected by plating the transformation mixture on LB agar (Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.) which had been supplemented with 50 mg/l of ampicillin.

Plasmid DNA was isolated from a transformant and checked by treatment with the restriction enzyme BamHI followed by agarose gel electrophoresis. The plasmid was called pUC18lyseE.

5 Example 2

Preparation of the dapB gene

Chromosomal DNA was isolated from the *Corynebacterium glutamicum* strain ATCC13032 as indicated in Example 1. The sequence of the dapB gene as such from *Corynebacterium glutamicum* is known (accession number X67737). However, the published DNA sequence comprises only 56 bp upstream from the translation start, so the 5' end upstream from the translation start was additionally sequenced.

The sequencing was carried out with plasmid pJC25 (EP-B 0 435 132) using a primer oligonucleotide which binds in the region of the known dapB sequence (accession number X67737). The sequence of the sequencing primer used was:

5' GAA CGC CAA CCT TGA TTC C 3'

The sequencing was carried out by the chain termination method described by Sanger et al., Proc. Natl. Acad. Sci. USA, (74) 5463-5467 (1977). The sequencing reaction was performed with the aid of the AutoRead Sequencing Kit (Pharmacia, Freiburg). The electrophoretic analysis and detection of the sequencing products were carried out with the A.L.F. DNA sequencer from Pharmacia (Freiburg, Germany).

The DNA sequence obtained was used to choose a second primer in order to obtain further sequence data upstream from the transcription start. The following primer was chosen for this purpose:

5' CTT TGC CGC CGT TGG GTT C 3'

The sequencing reaction was carried out as described above. The novel sequence upstream from the dapB gene is shown as SEQ ID No. 1. The sequence including the nucleotide sequence of the dapB gene is shown as SEQ ID

5 No. 2.

The polymerase chain reaction was used to amplify the dapB gene. For this purpose, two primer oligonucleotides, chosen on the basis of the known DNA sequence of the dapB gene, were synthesized by MWG Biotech:

10 P-dap:

5' (AAG CTT) AGG TTG TAG GCG TTG AGC 3'

dapall:

5' TTA ACT TGT TCG GCC ACA GC 3'

The 5' primer (primer P-dap) contains a HindIII cleavage site which is indicated by brackets in the sequence shown above. The PCR was carried out as in Example 1. An approx. 1.1 kb DNA fragment, which carries the dapB gene and contains a cleavage site for the restriction endonuclease HindIII at each end, was amplified in this way. The PCR fragment obtained was purified from 0.8% agarose gel (QIAquick Gel Extraction Kit from Qiagen, Hilden, Germany) and cloned into cloning vector pCR2.1TOPO (Invitrogen, Leek, The Netherlands) with the TOPO TA Cloning Kit (Invitrogen, Leek, The Netherlands, cat. no. K4550-01). The ligation mixture was transformed to the E. coli strain TOP10F' from Invitrogen, the transformation mixture was plated on LB agar containing kanamycin (50 mg/l), IPTG (0.16 mM) and X-Gal (64 mg/l) and kanamycin-resistant, white colonies were isolated. Plasmid DNA was isolated from a transformant with the aid of the QIAprep Spin Miniprep Kit from Qiagen and checked by cleavage with the restriction enzyme HindIII followed by agarose gel electrophoresis. The DNA sequence of the amplified DNA fragment was checked by sequencing. The sequence of the

PCR product matches the sequence shown in SEQ ID No. 1.
The plasmid obtained was called pCR2.1TOPOdapB.

Example 3

Preparation of the DNA coding for the *pyc* gene

5 The *Corynebacterium glutamicum* strain ATCC13032 was used
as the donor for the chromosomal DNA. Chromosomal DNA was
isolated from the strain ATCC13032 as described in Example
1. The polymerase chain reaction was used to amplify a
DNA fragment carrying the *pyc* gene. The following primer
10 oligonucleotides were chosen for the PCR on the basis of
the *pyc* gene sequence known for *C. glutamicum* (Peters-
Wendisch et al., Microbiology 144, 915-927 (1998))
(accession number Y09548):

5-PYC-IN:

15 5' GC(T CTA GA)A GTG TCG CAA CCG TGC TTG A 3'

3-PYC-IN:

5' GC(T CTA GA)T TGA GCC TTG GTC TCC ATC T 3'

The primers shown were synthesized by MWG Biotech and the
PCR reaction was carried out by the standard PCR method of
20 Innis et al. (PCR Protocols. A Guide to Methods and
Applications, 1990, Academic Press). The primers make it
possible to amplify an approx. 3.8 kb DNA fragment
carrying the *pyc* gene. The primers also contain the
sequence for a cleavage site of the restriction
25 endonuclease *Xba*I, which is indicated by brackets in the
nucleotide sequence shown above.

The amplified DNA fragment of approx. 3.8 kb, carrying the
pyc gene, was identified by gel electrophoresis in 0.8%
agarose gel, isolated from the gel and purified by the
30 conventional methods (QIAquick Gel Extraction Kit, Qiagen,
Hilden).

The fragment was then ligated to vector pCRII-TOPO by means of the Dual Promotor Topo TA Cloning Kit (Invitrogen, Leek, The Netherlands, cat. number K4600-01). The ligation mixture was transformed to the E. coli strain TOP10 (Invitrogen, Leek, The Netherlands). Plasmid-carrying cells were selected by plating the transformation mixture on LB agar containing kanamycin (50 mg/l) and X-Gal (64 mg/l).

After isolation of the DNA, the plasmid obtained was checked by means of restriction cleavage and identified in agarose gel. The plasmid was called pCRII-TOPO_{pyc} and the DNA sequence of the cloned insert was sequenced for control purposes. As the determined sequence of the pyc insert in pCRII-TOPO_{pyc} matches the sequence of the gene library entry, this plasmid was used subsequently.

Example 4

Cloning of the dapB gene into vector pEC7

An approx. 1.1 kb DNA fragment carrying the dapB gene was isolated from plasmid pCR2.1TOPO_{dapB} (from Example 2).

For this purpose, plasmid pCR2.1TOPO_{dapB} was fully digested with the restriction enzyme HindIII and the approx. 1.1 kb DNA fragment carrying the dapB gene was isolated.

The dapB fragment was inserted into vector pEC7. Vector pEC7 is based on E. coli - C. glutamicum shuttle vector pEC5 (Eikmanns et al., 1991: 93-98). The BamHI cleavage site not located in the polylinker was removed from plasmid pEC5 in the following manner: Plasmid pEC5 was partially cleaved with the restriction enzyme BamHI.

The approx. 7.2 kb DNA fragment was isolated from the agarose gel and the protruding ends were filled in with Klenow polymerase (Boehringer Mannheim). The resulting DNA fragment was ligated (T4 ligase, Boehringer Mannheim). The ligation mixture was transformed to the E. coli strain

DH5 α and kanamycin-resistant colonies were isolated on LB agar containing kanamycin (50 mg/l). Plasmid DNA was isolated from a transformant (QIAprep Spin Miniprep Kit from Qiagen) and checked by restriction cleavage with the
5 restriction enzymes BamHI and PstI. The resulting plasmid was called pEC6.

Plasmid pEC6 was fully cleaved with the restriction enzyme XhoI. A DNA fragment carrying the trp terminator was ligated to vector DNA fragment (T4 ligase, Boehringer
10 Mannheim). The ligation mixture was transformed to the E. coli strain DH5 α and kanamycin-resistant colonies were isolated on LB agar containing kanamycin (50 mg/l). Plasmid DNA was isolated from a transformant (QIAprep Spin Miniprep Kit from Qiagen) and checked by restriction
15 cleavage with the restriction enzymes BamHI and XhoI. The resulting plasmid was called pEC7.

The dapB-carrying DNA fragment obtained was ligated to vector pEC7 (T4 DNA ligase, Boehringer Mannheim), which had also been fully digested with the restriction enzyme
20 HindIII and treated with alkaline phosphatase (Boehringer Mannheim). The ligation mixture was transformed to the E. coli strain DH5 α and kanamycin-resistant colonies were isolated on LB agar containing kanamycin (50 mg/l). Plasmid DNA was isolated from a transformant (QIAprep Spin
25 Miniprep Kit from Qiagen) and checked by restriction cleavage with the restriction enzyme HindIII. The resulting plasmid was called pEC7dapB (Figure 1). The Escherichia coli strain obtained was called DH5 α /pEC7dapB.

Example 5

30 Cloning of the lysE gene into vector pEC7

Plasmid pUC18lysEneu described in Example 1 was fully digested with the restriction enzyme BamHI and the 1.1 kb BamHI fragment carrying the lysE gene was isolated as in Example 1. Vector pEC7 was likewise fully cleaved with

the restriction enzyme BamHI and treated with alkaline phosphatase. The BamHI vector fragment and the BamHI lysE fragment were ligated (Rapid DNA Ligation Kit, Boehringer Mannheim) and transformed to the E. coli strain DH5 α .

- 5 Plasmid-carrying transformants were selected on LB agar containing chloramphenicol (10 mg/l). Plasmid DNA was isolated (QIAprep Spin Miniprep Kit, Qiagen) and checked by restriction cleavage with the enzyme BamHI. The resulting plasmid was called pEC7lysE (Figure 2). The
10 strain obtained by transformation of plasmid pEC7lysE to the E. coli strain DH5 α was called DH5 α /pEC7lysE.

Example 6

Cloning of the pyc gene into vector pEC7

- The 3.8 kb DNA fragment carrying the pyc gene from C.
15 glutamicum ATCC13032 was obtained from plasmid pCRII-TOPOpyc (from Example 3) by cleavage with the restriction enzyme XbaI. The 3.8 kb DNA fragment was identified by gel electrophoresis, isolated from the gel and purified by the conventional methods and the protruding ends were
20 filled in with Klenow polymerase. Vector pEC7 was likewise fully cleaved with the restriction enzyme SmaI and treated with alkaline phosphatase. The SmaI vector fragment and the XbaI pyc fragment treated with Klenow polymerase were ligated (T4 ligase, Boehringer Mannheim)
25 and transformed to the E. coli strain DH5 α . Plasmid-carrying transformants were selected on LB agar containing chloramphenicol (10 mg/l). Plasmid DNA was isolated (QIAprep Spin Miniprep Kit, Qiagen, Hilden, Germany) and checked by restriction cleavage with the restriction
30 enzyme SalI. The resulting plasmid was called pEC7pyc (Figure 3). The E. coli strain obtained by transformation of plasmid pEC7pyc to the E. coli strain DH5 α was called DH5 α /pEC7pyc.

Example 7

Preparation of a plasmid containing *lysE* and *dapB* genes

The *dapB* gene was isolated as a *Hind*III fragment from plasmid pCR2.1TOP0dapB containing the *dapB* gene from *C. glutamicum* ATCC13032. To do this, the plasmid was fully digested with the restriction enzyme *Hind*III and the *dapB*-carrying DNA fragment was isolated from 0.8% agarose gel (QIAquick Gel Extraction Kit, Qiagen).

Vector pEC7lysE was also fully digested with the restriction enzyme *Hind*III and treated with alkaline phosphatase. The 1.1 kb fragment containing *dapB* was ligated to the resulting linear vector fragment (T4 ligase, Boehringer Mannheim) and the ligation mixture was transformed to the *E. coli* strain DH5 α . Plasmid-carrying transformants were selected on LB agar containing chloramphenicol (10 mg/l). Plasmid DNA was isolated (QIAprep Spin Miniprep Kit, Qiagen, Hilden, Germany) and checked by restriction cleavage with the restriction enzyme *Hind*III.

The resulting plasmid was called pEC7lysEdapB. This plasmid is capable of autonomous replication in *Escherichia coli* and in *Corynebacterium* and confers resistance to the antibiotic chloramphenicol on its host.

Plasmid pEC7lysEdapB simultaneously contains the *dapB* gene, which codes for dihydrodipicolinate reductase, and the *lysE* gene, which codes for the lysine exporter.

The strain obtained by the transformation of *E. coli* DH5 α with pEC7lysEdapB was called DH5 α /pEC7lysEdapB.

Example 8

Preparation of a plasmid simultaneously containing dapB and pyc genes

The plasmid carrying the pyc gene which codes for the pyruvate carboxylase from *Corynebacterium glutamicum* ATCC13032 was fully cleaved with the restriction enzyme XbaI and the protruding ends were filled in with Klenow polymerase as described in Example 6, making it possible to isolate the 3.8 kb DNA fragment containing the gene for pyruvate carboxylase.

Plasmid pEC7dapB (from Example 4) was also fully cleaved with the restriction enzyme SmaI and the ends were treated with alkaline phosphatase. The resulting linear vector fragment was ligated to the 3.8 kb DNA fragment containing the pyc gene using T4 DNA ligase (Boehringer Mannheim, Mannheim, Germany). This produced a plasmid containing both the dapB gene and the pyc gene from *corynebacteria*. Plasmid-carrying transformants were selected on LB agar containing chloramphenicol (10 mg/l). Plasmid DNA was isolated (QIAprep Spin Miniprep Kit, Qiagen, Hilden, Germany) and verified by restriction cleavage with the restriction enzyme SalI. The plasmid is shown in Figure 4 and was called pEC7dapBpyc. The *E. coli* strain obtained by transformation of plasmid pEC7dapBpyc to the *E. coli* strain DH5 α was called DH5 α /pEC7dapBpyc.

Example 9

Preparation of a plasmid containing sequences simultaneously coding for the lyse, dapB and pyc genes

Plasmid pCRII-TOPOpyc (from Example 3), which carries the pyc gene coding for the pyruvate carboxylase from *Corynebacterium glutamicum* ATCC13032, was fully cleaved with the restriction enzyme XbaI and treated with Klenow polymerase as described in Example 6, making it possible

to isolate the 3.8 kb DNA fragment containing the gene for pyruvate carboxylase.

Plasmid pEC7dapBlysE was also fully cleaved with the restriction enzyme SmaI and the ends were treated with alkaline phosphatase. The resulting linear vector fragment was ligated to the 3.8 kb DNA fragment containing the pyc gene using T4 DNA ligase (Boehringer Mannheim). This produces a plasmid containing the lysE gene and dapB gene and the pyc gene from *Corynebacterium glutamicum*.

Plasmid-carrying transformants were selected on LB agar containing chloramphenicol (10 mg/l). Plasmid DNA was isolated (QIAprep Spin Miniprep Kit, Qiagen, Hilden, Germany) and verified by restriction cleavage with the restriction enzyme SmaI. The plasmid is shown in Figure 5 and was called pEC7lysEdapBpyc. The *E. coli* strain obtained by transformation of plasmid pEC7dapBlysEpyc to the *E. coli* strain DH5 α was called DH5 α /pEC7dapBlysEpyc.

Example 10

Transformation of the strain MH20-22B with plasmids pJC23 and pJC33

Plasmid pJC1 is a plasmid capable of replication in *Escherichia coli* and *Corynebacterium glutamicum* (Cremer et al., Molecular and General Genetics 220: 478-480 (1990)). Plasmid pJC33 (Cremer et al., Applied and Environmental Microbiology 57(6), 1746-1752 (1991)), which carries the lysC(Fbr) gene from the *C. glutamicum* strain MH20-22B, is derived therefrom.

Plasmid pJC23 is also based on vector pJC1 and carries the dapA gene from *C. glutamicum* ATCC13032 (Cremer et al., Molecular and General Genetics 220: 478-480 (1990)) (EP-B 0 435 132). Plasmids pJC1, pJC33 and pJC23 were introduced into the strain MH20-22B by the electroporation method (Haynes and Britz, FEMS Microbiology Letters (61) 329-334 (1989)). The *C. glutamicum* strain MH20-22B is an

AEC-resistant lysine producer deposited under the number DSM5715.

The transformants obtained by means of electroporation were isolated on selection agar (LBHIS agar (18.5 g/l of brain-heart infusion broth, 0.5 M sorbitol, 5 g/l of bacto tryptone, 2.5 g/l of bacto yeast extract, 5 g/l of NaCl, 18 g/l of bacto agar)) containing 15 mg/l of kanamycin. Plasmid DNA was isolated by the conventional methods (Peters-Wendisch et al., Microbiology 144, 915-927 (1998)), cleaved with suitable restriction endonucleases and checked. The strains obtained were called MH20-22B/pJC1, MH20-22B/pJC33 and MH20-22B/pJC23.

Example 11

Transformation with plasmids pEC7pyc, pEC7dapBpyc and pEC7lysEdapBpyc

The strains prepared in Example 10 were subsequently provided with a second plasmid.

The following plasmids were introduced by the electroporation method into the strains MH20-22B/pJC1, MH20-22B/pJC33 and MH20-22B/pJC23 described:

- pEC7pyc (cf. Example 6)
- pEC7dapBpyc (cf. Example 8)
- pEC7lysEdapBpyc (cf. Example 9)

The transformed bacteria are selected on the basis of the antibiotic resistance of the plasmids they contain. The transformants obtained by means of electroporation were isolated on selection agar (LBHIS agar containing 15 mg/l of kanamycin and 7.5 mg/l of chloramphenicol). Plasmid DNA was isolated, cleaved with suitable restriction endonucleases and checked.

The strains obtained are listed below:

- DSM5715/pJC1/pEC7pyc
- DSM5715/pJC33/pEC7pyc
- DSM5715/pJC23/pEC7pyc
- 5 DSM5715/pJC23/pEC7dapBpyc
- DSM5715/pJC23/pEC7lysEdapBpyc

Example 12

Preparation of L-lysine

- 10 The various *C. glutamicum* strains obtained in Examples 10 and 11 were cultivated in a nutrient medium suitable for lysine production and the lysine content of the culture supernatant was determined.

- 15 This was done by first incubating the various strains on agar plates with the appropriate antibiotics (brain-heart agar containing kanamycin (25 mg/l) and chloramphenicol (10 mg/l)) for 24 hours at 33°C. These agar plate cultures were used to inoculate a preculture (10 ml of medium in a 100 ml conical flask). Complete medium CgIII was used as the preculture medium. Kanamycin (25 mg/l) and
- 20 chloramphenicol (10 mg/l) were added. The preculture was incubated for 24 hours at 33°C on a shaker at 250 rpm. This preculture was used to inoculate a main culture to give an initial OD (660 nm) of 0.2. Medium MM was used for the main culture.

Medium MM

CSL 5 g/l
MOPS 20 g/l
Glucose 50 g/l (autoclave separately)

5 Salts:

(NH₄)₂SO₄ 25 g/l
KH₂PO₄ 0.1 g/l
MgSO₄*7H₂O 1.0 g/l
CaCl₂*2H₂O 10 mg/l
10 FeSO₄*7H₂O 10 mg/l
MnSO₄*H₂O 5.0 mg/l
Biotin 0.3 mg/l (sterile-filtered)
Thiamine*HCl 0.2 mg/l (sterile-filtered)
CaCO₃ 25 g/l

15 Abbreviations:

CSL: corn steep liquor
MOPS: morpholinopropanesulfonic acid

CSL, MOPS and the salt solution are adjusted to pH 7 with aqueous ammonia and autoclaved. The sterile substrate and
20 vitamin solutions and the dry-autoclaved CaCO₃ are then added.

Cultivation is carried out in a volume of 10 ml in a 100 ml conical flask with baffles. Kanamycin (25 mg/l) and chloramphenicol (10 mg/l) were added. Cultivation
25 proceeded at 33°C and 80% atmospheric humidity.

After 72 hours the OD was measured at a wavelength of 660 nm. The amount of lysine formed was determined with an amino acid analyzer from Eppendorf BioTronik (Hamburg, Germany) by means of ion exchange chromatography and
30 postcolumn derivation with ninhydrin detection. The glucose content was determined with a sugar analyzer from Skalar Analytik GmbH (Erkelenz, Germany).

The experimental results are shown in Table 1.

Table 1

Strain	Gene	OD (660 nm)	Lysine-HCl g/l
DSM5715/pJC1/pEC7pyc	pyc	9.3	11.3
DSM5715/pJC33/pEC7pyc	pyc, lysC(Fbr)	9.2	11.9
DSM5715/pJC23/pEC7pyc	pyc, dapA	9.3	14.4
DSM5715/pJC23/pEC7dapBpyc	pyc, dapA, dapB	9.3	16.9
DSM5715/pJC23/ pEC7lysEdapBpyc	pyc, dapA, dapB, lysE	9.1	17.6

Example 13

5 Cloning of the aecD gene into vector pUC18

Plasmid pSIR21 (Rossol, Thesis, University of Bielefeld 1992) was fully cleaved with the enzymes BglII and EcoRV and the 1.4 kb DNA fragment containing the aecD gene (accession number M89931) (Rossol and Pühler, Journal of Bacteriology 174 (9), 2968-2977 (1992)) from *C. glutamicum* ATCC13032 was isolated. The isolated DNA fragment was ligated to plasmid pUC18 (which had been fully digested with the enzymes BamHI and SmaI) using T4 DNA ligase, as described in Sambrook et al. (Molecular Cloning: A Laboratory Manual (1989), Cold Spring Harbor Laboratory Press). The ligation mixture was transformed to the *E. coli* strain DH5 α . The transformants were selected on brain-heart agar plates containing 100 mg/l of ampicillin. Plasmid DNA was isolated from one colony. The plasmid obtained was called pUC18::aecD.

Example 14

Cloning of the dapA gene into plasmid pSP72

A dapA gene fragment is isolated from plasmid pJC20 (Cremer, J., Thesis 1989, University of Düsseldorf) as an SphI-BamHI fragment. Vector pSP72 (Promega Corporation, USA) was fully cleaved with the enzymes SphI and BamHI and treated with alkaline phosphatase. The dapA-carrying fragment was ligated to this vector using T4 DNA ligase. The DNA was then transformed to the E. coli strain XL1 Blue (Bullock, Fernandez and Short, BioTechniques (5), 376-379 (1987)). The transformants were selected on LB medium containing 100 mg/l of ampicillin. Plasmid DNA was isolated from one transformant and called pSP72::dapA.

Example 15

15 Mutagenesis of the dapA promoter and preparation of plasmids pSP72::dapA(MC20) and pSP72::dapA(MA16)

The Quickchange site directed mutagenesis kit from Stratagene was used for the mutagenesis of the promoter region. The following primers were constructed with the aid of said dapA sequence and used for the mutagenesis:

For the preparation of pSP72::dapA(MC20)

Primer dap1 for MC20

CCA AAT GAG AGA TGG TAA CCT TGA ACT CTA TGA GCA

Primer dap2 for MC20

25 GTG CTC ATA GAG TTC AAG GTT ACC ATC TTC CCT CAT TTG G

For the preparation of pSP72::dapA(MA16)

Primer dap3 for MA16

CCA AAT GAG GGA AGA AGG TAT AAT TGA ACT CTA TGA GCA

Primer dap4 for MA16

30 GTG CTC ATA GAG TTC AAT TAT ACC TTC TTC CCT CAT TTG G

The PCR was carried out as indicated by the manufacturer of the Quickchange site directed mutagenesis kit (Stratagene) using plasmid pSP72::dapA (from Example 14) as the template.

- 5 The mutagenesis mixtures were transformed to the E. coli strain XL1 Blue. The transformants were selected on LB medium containing 100 mg/l of carbenicillin. Plasmid DNA was isolated from one transformant and the loss of the BstEII cleavage site was controlled by BstEII digestion.
- 10 Plasmids no longer carrying a BstEII cleavage site exhibited the desired mutation.

- The plasmids obtained were transformed to the dapA-defective E. coli mutant RDA8. The transformation mixtures were plated on LB containing 100 mg/l of
- 15 carbenicillin in order to test the complementation of the dapA mutation. DNA was isolated from one transformant in each case and the plasmids obtained were called pSP72::dapA(MC20) and pSP72::dapA(MA16). The plasmids were sequenced by the chain termination method described
- 20 in Sanger et al., Proceedings of the National Academy of Sciences of the USA (74), 5463-5467 (1977), using the reverse and universal sequencing primers. The sequencing reaction was performed with the aid of the AutoRead Sequencing Kit (Pharmacia, Freiburg). The electrophoretic
- 25 analysis and detection of the sequencing products were carried out with the A.L.F. DNA sequencer (Pharmacia, Freiburg, Germany).

Example 16

- Preparation of plasmids pK19mobsacBaecD::dapA(MC20) and
- 30 pK19mobsacBaecD::dapA(MA16) (recloning of the mutagenized fragments)

Plasmids pSP72::dapA(MC20) and pSP72::dapA(MA16) (from Example 15) were fully cleaved with the restriction enzymes PvuII and SmaI. The 1450 bp PvuII-SmaI fragments

carrying the *dapA* gene with the mutated MC20 or MA16 promoter were ligated to *Stu*I-cleaved vector pUC18::*aecD* (from Example 13) using T4 DNA ligase. The ligation mixture was transformed to the *E. coli* strain DH5 α . The transformants were selected on LB medium containing 100 mg/l of ampicillin. Plasmid DNA was isolated from one transformant in each case to give plasmids pUC18*aecD*::*dapA*(MC20) and pUC18*aecD*::*dapA*(MA16).

Plasmids pUC18*aecD*::*dapA*(MC20) and pUC18*aecD*::*dapA*(MA16) were partially cleaved with the restriction enzyme *Eco*RI and fully cleaved with the enzyme *Sal*I to give the 3.0 kb fragment carrying *aecD*::*dapA*(MA16) or *aecD*::*dapA*(MC20). The fragment was ligated to vector pK19*mobsacB* (which had been cleaved and treated with alkaline phosphatase) (Schäfer et al., Gene (145), 69-73 (1994)) using T4 DNA ligase. The ligation mixture was transformed to the *E. coli* strain DH5 (Hanahan (1985), in: DNA Cloning. A Practical Approach, Vol. I, IRL-Press, Oxford, Washington DC, USA). The transformants were selected on LB medium containing 50 mg/l of kanamycin. Plasmid DNA was isolated from one transformant in each case to give plasmids pK19*mobsacBaecD*::*dapA*(MC20) and pK19*mobsacBaecD*::*dapA*(MA16).

The plasmid DNA was transformed to the *E. coli* strain S17-1 (Simon, Priefer and Pühler, Bio/Technology (1), 784-791 (1983)). The transformants were selected on LB medium containing 50 mg/l of kanamycin. Plasmid DNA was isolated from one transformant in each case and checked. The strains obtained were called S17-1/pK19*mobsacBaecD*::*dapA*(MC20) and S17-1/pK19*mobsacBaecD*::*dapA*(MA16).

Example 17

Preparation of the *C. glutamicum* strains

DSM5715aecD::dapA(MC20) and DSM5715aecD::dapA(MA16)

Plasmids pK19mobsacBaecD::dapA(MC20) and

5 pK19mobsacBaecD::dapA(MA16) were transferred from

S17-1/pK19mobsacBaecD::dapA(MC20) and

S17-1/pK19mobsacBaecD::dapA(MA16) (from Example 16) to the
C. glutamicum strain DSM5715 by the conjugation method

(Schäfer et al., Journal of Bacteriology (172), 1663-1666

10 (1990)). For selection of the transconjugants, the

conjugation mixtures were plated on brain-heart medium
containing nalidixic acid and kanamycin. The

transconjugants obtained were incubated overnight in 10 ml
of brain-heart medium. Aliquots were then plated on

15 plates containing sucrose (brain-heart agar containing 10%
of sucrose) in order to select for loss of sucrose

sensitivity. Sucrose-resistant clones were isolated and
checked again on agar plates containing chloramphenicol

and kanamycin (brain-heart medium containing 15 mg/l of

20 kanamycin and brain-heart medium containing 10 mg/l of
chloramphenicol).

Colonies exhibiting the following phenotype were isolated:

sucrose resistant

kanamycin sensitive

25 chloramphenicol sensitive

The insertion of the dapA gene fragment into the aecD gene
was checked by the Southern blot method (Sambrook et al.,
Molecular Cloning: A Laboratory Manual (1989), Cold Spring
Harbor Laboratory Press).

Example 18

Preparation of the *C. glutamicum* strains

DSM5715aecD::dapA(MC20)/pEC7pyc,

DSM5715aecD::dapA(MA16)/pEC7pyc,

- 5 DSM5715aecD::dapA(MC20)/pEC7, DSM5715aecD::dapA(MA16)/pEC7
and DSM5715/pEC7

As described in Example 6, the pyc gene is present in vector pEC7pyc. This plasmid pEC7pyc and plasmid pEC7 were introduced into the strains DSM5715aecD::dapA(MC20),

- 10 DSM5715aecD::dapA(MA16) and DSM5715 (from Example 17) by means of electroporation (Haynes 1989, FEMS Microbiology Letters 61, 329-334) to give *C. glutamicum*
DSM5715aecD::dapA(MC20)/pEC7pyc,
DSM5715aecD::dapA(MA16)/pEC7pyc,
15 DSM5715aecD::dapA(MC20)/pEC7, DSM5715aecD::dapA(MA16)/pEC7
and DSM5715/pEC7. The transformants were selected on brain-heart agar containing 25 mg/l of kanamycin. Plasmid DNA was isolated from one transformant in each case and checked.

- 20 The following strains were obtained in this way:

DSM5715aecD::dapA(MC20)/pEC7pyc,

DSM5715aecD::dapA(MA16)/pEC7pyc,

DSM5715aecD::dapA(MC20)/pEC7,

DSM5715aecD::dapA(MA16)/pEC7 and

- 25 DSM5715/pEC7.

Example 19

Preparation of lysine with the strains prepared in Example 18

- After precultivation in CgIII medium (Kase & Nakayama, Agricultural and Biological Chemistry 36 (9), 1611-1621 (1972)), the strains DSM5715aecD::dapA(MC20)/pEC7pyc,
- 30

DSM5715aecD::dapA(MA16)/pEC7pyc,
DSM5715aecD::dapA(MC20)/pEC7, DSM5715aecD::dapA(MA16)/pEC7
and DSM5715/pEC7 were cultivated in MM production medium
as described in Example 12. After incubation for 48
5 hours, the optical density at 660 nm and the concentration
of L-lysine formed were determined.

The experimental results are shown in Table 2.

Table 2

Strain	OD (660 nm)	Lysine·HCl g/l
DSM5715aecD::dapA(MC20)/pEC7pyc	14.15	13.3
DSM5715aecD::dapA(MA16)/pEC7pyc	13.6	13.4
DSM5715/pEC7	13.3	11.5
DSM5715aecD::dapA(MA16)/pEC7	13.3	12.9
DSM5715aecD::dapA(MC20)/pEC7	14.0	12.8

10 Example 20

Transformation of strain DM678 with plasmids pEC7 and
pEC7lysEdapBpyc

The lysine producing *C. glutamicum* strain DM678 was
derived from ATCC13032 by several rounds of mutagenesis
15 and screening. It has a partial requirement for L-
threonine and its growth is sensitive towards L-
methionine. Strain DM678 is deposited as DSM12866 at the
Deutsche Sammlung für Mikroorganismen und Zellkulturen
(DSMZ, Braunschweig, Deutschland).

Plasmids pEC7 (from Example 4) and pEC7lysEdapBpyc (from Example 9) were introduced into the strain DM678 by the electroporation method (Haynes and Britz, FEMS Microbiology Letters (61) 329-334 (1989)).

- 5 The transformants obtained by means of electroporation were isolated on selection agar (LBHIS agar (18.5 g/l of brain-heart infusion broth, 0.5 M sorbitol, 5 g/l of bacto tryptone, 2.5 g/l of bacto yeast extract, 5 g/l of NaCl, 18 g/l of bacto agar)) containing 7.5 mg/l of
10 chloramphenicol. Plasmid DNA was isolated by the conventional methods (Peters-Wendisch et al., Microbiology 144, 915-927 (1998)), cleaved with suitable restriction endonucleases and checked by agarosegel electrophoresis. The strains obtained were called DM678/pEC7 and
15 DM678/pEC7lysEdapBpyc.

Example 21

Preparation of L-lysine

- The *C. glutamicum* strains obtained in Examples 20 were cultivated in a nutrient medium suitable for lysine
20 production and the lysine content of the culture supernatant was determined.

- This was done by first incubating the strains on agar plates (brain-heart) with chloramphenicol (10 mg/l) for 24 hours at 33°C. These agar plate cultures were used to
25 inoculate a preculture (10 ml of medium in a 100 ml conical flask). Complete medium CgIII which had been supplemented with 10 mg/l of chloramphenicol was used as the preculture medium. The preculture was incubated for 24 hours at 33°C on a shaker at 250 rpm. This preculture was
30 used to inoculate a main culture to give an initial OD (660 nm) of 0.1. Medium MM2 was used for the main culture.

Medium MM2:

CSL	25 g/l
MOPS	20 g/l
Glucose	80 g/l (autoclave separately)

5 Salts:

(NH ₄) ₂ SO ₄	25 g/l
KH ₂ PO ₄	0.1 g/l
MgSO ₄ *7H ₂ O	0.25 g/l
CaCl ₂ *2H ₂ O	10 mg/l
10 FeSO ₄ *7H ₂ O	10 mg/l
MnSO ₄ *H ₂ O	10 mg/l
Biotin	0.3 mg/l (sterile-filtered)
Thiamine*HCl	0.2 mg/l (sterile-filtered)
CaCO ₃	25 g/l (autoclave separately)

15 Abbreviations:

CSL: corn steep liquor
MOPS: morpholinopropanesulfonic acid

CSL, MOPS and the salt solution are adjusted to pH 7 with aqueous ammonia and autoclaved. The sterile substrate and vitamin solutions, the Chloramphenicol (10 mg/l) and the dry-autoclaved CaCO₃ are then added.

Cultivation is carried out in a volume of 10 ml in a 100 ml conical flask with baffles. Incubation proceeded at 33°C and 80% atmospheric humidity.

25 After 48 hours the OD was measured at a wavelength of 660 nm. The amount of lysine formed was determined with an amino acid analyzer from Eppendorf BioTronik (Hamburg, Germany) by means of ion exchange chromatography and postcolumn derivation with ninhydrin detection. The
30 glucose content was determined with a sugar analyzer from Skalar Analytik GmbH (Erkelenz, Germany).

The experimental results are shown in Table 3.

Table 3

Strain	Gene	OD (660 nm)	Lysine-HCl g/l
DM678/pEC7		25.6	16.7
DM678/ pEC7lysEdapBpyc	pyc, dapB, lysE	23.3	17.7

Brief Description of the Figures

Figure 1 illustrates the pEC7dapB plasmid (cf. Example 4)

5 Figure 2 illustrates the pEC7lysE plasmid (cf. Example 5)

Figure 3 illustrates the pEC7pyc plasmid (cf. Example 6)

Figure 4 illustrates the pEC7dapBpyc plasmid (cf. Example 8)

10 Figure 5 illustrates the pEC7lysEdapBpyc plasmid (cf. Example 9).

The abbreviations used in the Figures are defined as follows:

Cm: chloramphenicol resistance gene

dapB: dapB gene from *C. glutamicum*

15 lysE: lysE gene from *C. glutamicum*

pyc: pyc gene from *C. glutamicum*

OriE: plasmid-coded origin of replication of *E. coli*

pBL: DNA fragment of plasmid pBL1

EcoRI: cleavage site of the restriction enzyme EcoRI

20 EcoRV: cleavage site of the restriction enzyme EcoRV

HincII: cleavage site of the restriction enzyme HincII

HindIII: cleavage site of the restriction enzyme HindIII

KpnI: cleavage site of the restriction enzyme KpnI

SalI: cleavage site of the restriction enzyme SalI

SmaI: cleavage site of the restriction enzyme SmaI

5 SphI: cleavage site of the restriction enzyme SphI

PvuII: cleavage site of the restriction enzyme PvuII

BamHI: cleavage site of the restriction enzyme BamHI

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
217